

The universal quantum invariant and colored ideal triangulations

Sakie Suzuki

RIMS/The Hakubi Center, Kyoto university

2017.12.18 @ Strasbourg University

Introduction

Drinfeld double and Heisenberg double

Universal quantum invariant and its reconstruction

Extension

3-dim. descriptions

Introduction

- ▶ Background
- ▶ Ideas for reconstruction of quantum invariants
- ▶ Remarks & Related topics

Background

1984 Jones polynomial




“Quantum invariants”

- ▶ Colored Jones polynomial
- ▶ Reshetkhin–Turaev invariant
- ▶ Universal quantum invariant
- ▶ Kontsevich integral

Background

KEY POINT FOR CONSTRUCTIONS



“ R -matrix”

RIII move



“hexagon identity”

Background

- ▶ Reshetkhin-Turaev invariant

$R \in \text{End}(V \otimes V)$, V : fin.dim. linear sp.

$$(1 \otimes R)(R \otimes 1)(1 \otimes R) = (R \otimes 1)(1 \otimes R)(R \otimes 1)$$


- ▶ Universal quantum invariant

$R \in \mathfrak{R}^{\otimes 2}$, \mathfrak{R} : ribbon Hopf algebra

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$$

Background

Definitions are **combinatorial** and **diagrammatic**



⇒ It is not easy to see **topological properties** of links from quantum invariants.

Background

What are “topological properties” of links?

- ▶ Properties defined using **simple operations** or **surfaces**. e.g. invertible, achiral, Brunnian, ribbon, boundary, etc.
- ▶ Properties defined by **classical invariants**. e.g. genus, homology, fundamental group, bridge number, Milnor invariants, etc.

Background

TASK

Find relationships between **quantum invariants**
and **topological properties** of links!

Background

METHODS

Link (3-dim. obj.)
(w/ topological properties)



Link diagram (2-dim. obj.) \rightsquigarrow Quantum invariants
(w/ planer properties)

Background

METHODS

Link (3-dim. obj.) \rightarrow triangulation (3-dim. obj.)
(w/ topological properties)



Link diagram (2-dim. obj.) \rightsquigarrow Quantum invariants
(w/ planer properties)

Ideas for reconstruction of quantum invariants

A : a fin-dim Hopf algebra/ k

1. Drinfeld double $D(A) \sim_k A^* \otimes A$

$\Rightarrow R \in D(A)^{\otimes 2}$ s.t.

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12} \in D(A)^{\otimes 3}.$$

2. Heisenberg double $H(A) \sim_k A^* \otimes A$

$\Rightarrow S \in H(A)^{\otimes 2}$ s.t.

$$S_{12}S_{13}S_{23} = S_{23}S_{12} \in H(A)^{\otimes 3}.$$

Ideas for reconstruction of quantum invariants

Theorem (Kashaev '97)

There is an algebra embedding

$$\phi: D(A) \rightarrow H(A) \otimes H(A)^{\text{op}},$$


s.t.

$$\phi^{\otimes 2}(R) = S''_{14} S_{13} \tilde{S}_{24} S'_{23}.$$

S', S'', \tilde{S} : modifications of S satisfying pentagon relations


Ideas for reconstruction of quantum invariants

Octahedral triangulations of link complements



Ideas for reconstruction of quantum invariants


Octahedral triangulations of link complements



$$\phi^{\otimes 2}(R) = S''_{14} S_{13} \tilde{S}_{24} S'_{23}$$


Ideas for reconstruction of quantum invariants

Pachner (2, 3) move



Ideas for reconstruction of quantum invariants

Pachner (2, 3) move




$$S_{23}S_{12} = S_{12}S_{13}S_{23}$$

Ideas for reconstruction of quantum invariants

TO SUM UP...

Idea for the reconstruction



\mapsto

S -tensor

Pachner (2,3) move


\mapsto

pentagon identity

Ideas for reconstruction of quantum invariants

TO SUM UP...

Idea for the reconstruction



\mapsto

S -tensor

Pachner (2,3) move

\mapsto

pentagon identity


In this talk: w/ universal quantum invariant

Remarks

Turaev-Viro's state sum invariant for (M, \mathcal{T}) :

$$Z(M) = w^{-\#\{\text{verteces}\}} \sum_{\lambda} w_{\lambda} \prod_T W(T; \lambda)$$

- ▶ \mathcal{T} : a triangulation of M
- ▶ λ : a color (giving an integer on each edge)
- ▶ T : a tetrahedron in \mathcal{T}
- ▶ $W(T; \lambda) \in \mathbb{C}$: the weight on T




Remarks

Turaev-Viro's state sum invariant for (M, \mathcal{T}) :

$$Z(M) = w^{-\#\{\text{verteces}\}} \sum_{\lambda} w_{\lambda} \prod_T W(T; \lambda)$$

- ▶ \mathcal{T} : a triangulation of M
- ▶ λ : a color (giving an integer on each edge)
- ▶ T : a tetrahedron in \mathcal{T}
- ▶ $W(T; \lambda) \in \mathbb{C}$: the weight on T satisfying a **pentagon identity**.



Remarks

1. [Turaev-Viro]
(triangulation, quantum $6j$ -symbol)

$$\begin{vmatrix} j_1 & j_2 & j_3 \\ i_1 & i_2 & i_3 \end{vmatrix} \begin{vmatrix} j_1 & j_2 & j_3 \\ k_1 & k_2 & k_3 \end{vmatrix} = \sum_n [n]_q \begin{vmatrix} i_1 & i_2 & j_3 \\ k_2 & k_1 & n \end{vmatrix} \begin{vmatrix} i_2 & i_3 & j_1 \\ k_3 & k_2 & n \end{vmatrix} \begin{vmatrix} i_3 & i_1 & j_2 \\ k_1 & k_3 & n \end{vmatrix}$$


2. [Baseilhac-Benedetti] QHI
(ideal triangulation, quantum dilogarithm)

$$\Psi(V)\Psi(U) = \Psi(U)\Psi(-UV)\Psi(V)$$

Remarks

3. The universal quantum invariant
(link diagram, the universal R -matrix)


$$R = \sum_{i \geq 0} \alpha_i \otimes \beta_i \in D(A)^{\otimes 2}$$

J :  $\mapsto \sum_{i,j \geq 0} \beta'_j \alpha_i u^{-1} \otimes \alpha'_j \beta_i u$

Remarks

3. The universal quantum invariant
(link diagram, the universal R -matrix)

$$R = \sum_{i \geq 0} \alpha_i \otimes \beta_i \in D(A)^{\otimes 2}$$


J : 

$$\mapsto \underbrace{\sum_{i,j \geq 0}}_{\text{state sum}} \underbrace{\beta'_j \alpha_i u^{-1} \otimes \alpha'_j \beta_i u}_{\text{products of weights}}$$

Remarks

3. The universal quantum invariant
(link diagram, the universal R -matrix)

$$R = \sum_{i \geq 0} \alpha_i \otimes \beta_i \in D(A)^{\otimes 2}$$

J :  \mapsto $\sum_{i,j \geq 0} \beta'_j \alpha_i u^{-1} \otimes \alpha'_j \beta_i u$

↑
↑
 state sum products of weights

The **orientation** of the link \Rightarrow The **order of products** of weights.

Remarks

4. **Reconstruction** of the universal quantum invariant
(**colored** ideal triangulation, the S -tensor)

$$J': \quad \begin{array}{c} \text{Diagram of a diamond-shaped ideal triangulation with colored arcs (blue and red) and arrows.} \end{array} \quad \mapsto \quad S = \sum_{i \geq 0} \alpha_i \otimes \beta_i \in H(A)^{\otimes 2}$$

Remarks


4. **Reconstruction** of the universal quantum invariant
(**colored** ideal triangulation, the S -tensor)

$$J': \quad \begin{array}{c} \text{Diagram of a diamond-shaped ideal triangulation with a blue link on the left and a red link on the right, both with arrows pointing outwards.} \end{array} \quad \mapsto \quad S = \sum_{i \geq 0} \alpha_i \otimes \beta_i \in H(A)^{\otimes 2}$$

- ▶ **invariant for “colored” 3-mfds**
(\exists a canonical choice of the color for a link \Rightarrow link inv.)
- ▶ **invariant for closed 3-mfds** if A is involutory


Related topics; w/ the reconstruction J'

1. To study **topological properties** of links using J' .



Related topics; w/ the reconstruction J'

- To study **topological properties** of links using J' .




- e.g. Obtain the **volume** of the link complement from J' !
 (=Restate the volume conjecture using J')

Related topics; w/ the reconstruction J'

2. v.s. Phys?

Related topics; w/ the reconstruction J'

2. v.s. Phys?



e.g. Give a TQFT construction of J' !

Related topics; w/ the reconstruction J'

3. v.s. Kontsevich invariant

$$Z(\mathcal{G}) \in A(\mathcal{O}) = \text{Span}_{\mathbb{C}} \langle \bigcirc, \bigcirc, \bigcirc, \dots \rangle / \text{HX STU AS}$$

$$R = \begin{array}{|c|} \hline \diagup \quad \diagdown \\ \hline \text{exp} \frac{1}{2} \\ \hline \diagdown \quad \diagup \\ \hline \end{array}, \quad \exists W_{\mathcal{G}} \text{ s.t. } J_{\mathcal{G}}(k) = W_{\mathcal{G}0} Z(k)$$

Related topics; w/ the reconstruction J'

3. v.s. Kontsevich invariant

$$Z(\mathcal{G}) \in A(\mathcal{O}) = \text{Span}_{\mathbb{C}} \langle \bigcirc, \bigcirc, \bigcirc, \dots \rangle / \begin{matrix} \text{HX} \\ \text{STU} \\ \text{AS} \end{matrix}$$

$$R = \boxed{\exp \frac{1}{2}} \quad , \quad \exists W_{\mathfrak{g}} \text{ s.t. } J_{\mathfrak{g}}(k) = W_{\mathfrak{g}} \circ Z(k)$$

Extend the reconstruction of the universal invariant to that of Kontsevich invariant!

Related topics; w/ J' for colored 3-mfds

1. v.s. WRT invariant (with $U_n(q)$)

$$M = S^3_L, \quad \tau_3(M) = \text{TR}_3(J_L)$$

suzyeny presentation

Related topics; w/ J' for colored 3-mfds

1. v.s. WRT invariant (with $U_n(q)$)

$$M = S^3 \setminus L, \quad \tau_3(M) = \text{TR}_3(J_L)$$

surjective presentation


 $J'(M)$
↕ ?
 $J''(S^3 \setminus L)$

Find direct relationship between J' and WRT invariant!

Related topics; w/ J' for colored 3-mfds

2. v.s. Turaev-Viro invariant, QHI, and Kuperberg invariant


← by triangulations.



Related topics; w/ J' for colored 3-mfds

2. v.s. Turaev-Viro invariant, QHI, and Kuperberg invariant

← by triangulations.




Find direct relationship between J' and these invariants!

Related topics; w/ J' for colored 3-mfds

2. Invariant using an associator
(working project with Anderson Vera)

$$\Phi \in \mathbb{A}(\downarrow\downarrow\downarrow) = \text{Span}_{\mathbb{C}} \langle \downarrow\downarrow\downarrow, \downarrow\downarrow\downarrow, \dots \rangle / \begin{matrix} \text{IHS} \\ \text{STU} \\ \text{AS} \end{matrix}$$

satisfies a "pentagon" relation.


$$[\text{Bar-Natan, Dancso, '12}] \quad Z(\text{triangle}) = \Phi$$


Related topics; w/ J' for colored 3-mfds

2. Invariant using an associator
(working project with Anderson Vera)

$$\Phi \in \mathbb{A}(\downarrow\downarrow\downarrow) = \text{Span}_{\mathbb{C}} \langle \downarrow\downarrow\downarrow, \downarrow\downarrow\downarrow, \dots \rangle / \begin{matrix} \text{IHS} \\ \text{STU} \\ \text{AS} \end{matrix}$$

satisfies a "pentagon" relation.

$$[\text{Bar-Natan, Dancso, '12}] \quad Z(\text{triangle}) = \Phi$$


Construct an invariant using an associator! then find direct relationship with LMO invariant!


Related topics; algebraic aspect

“Quantum group theory” for Heisenberg double

$U_h(\mathfrak{g})$
 $\left\{ \begin{array}{l} \bullet R\text{-matrix} \\ \bullet \text{crystal basis} \end{array} \right.$
← motivated
Knot theory

Related topics; algebraic aspect

“Quantum group theory” for Heisenberg double



Construct “quantum group theory” for Heisenberg double
(with S -tensor and crystal basis?)

Introduction

Drinfeld double and Heisenberg double

- ▶ Quasi-triangular Hopf algebra and Ribbon Hopf algebra
- ▶ Drinfeld double and Heisenberg double

Quasi-triangular Hopf algebra

Quasi-triangular Hopf algebra $(\mathfrak{R}, \eta, m, \varepsilon, \Delta, \gamma, R)$: Hopf algebra with the universal R -matrix $R \in \mathfrak{R}^{\otimes 2}$ such that


$$\begin{aligned}\Delta^{\text{op}}(x) &= R\Delta(x)R^{-1} \quad \text{for } x \in \mathfrak{R}, \\ (\Delta \otimes 1)(R) &= R_{13}R_{23}, \quad (1 \otimes \Delta)(R) = R_{13}R_{12}.\end{aligned}$$

Quasi-triangular Hopf algebra

Quasi-triangular Hopf algebra $(\mathfrak{A}, \eta, m, \varepsilon, \Delta, \gamma, R)$: Hopf algebra with the universal R -matrix $R \in \mathfrak{A}^{\otimes 2}$ such that

$$\begin{aligned}\Delta^{\text{op}}(x) &= R\Delta(x)R^{-1} \quad \text{for } x \in \mathfrak{A}, \\ (\Delta \otimes 1)(R) &= R_{13}R_{23}, \quad (1 \otimes \Delta)(R) = R_{13}R_{12}.\end{aligned}$$

\Rightarrow invariant for **braids**.



Ribbon Hopf algebra

Ribbon Hopf algebra $(\mathfrak{R}, \eta, m, \varepsilon, \Delta, \gamma, R, \theta)$: quasi-triangular Hopf algebra with the ribbon element $\theta \in \mathfrak{R}$ such that

$$\theta^2 = u\gamma(u), \quad \gamma(\theta) = \theta, \quad \varepsilon(\theta) = 1, \quad \Delta(\theta) = (R_{21}R)^{-1}(\theta \otimes \theta),$$

where $u = \sum \gamma(\beta)\alpha$ with $R = \sum \alpha \otimes \beta$.


Ribbon Hopf algebra

Ribbon Hopf algebra $(\mathfrak{R}, \eta, m, \varepsilon, \Delta, \gamma, R, \theta)$: quasi-triangular Hopf algebra with the ribbon element $\theta \in \mathfrak{R}$ such that

$$\theta^2 = u\gamma(u), \quad \gamma(\theta) = \theta, \quad \varepsilon(\theta) = 1, \quad \Delta(\theta) = (R_{21}R)^{-1}(\theta \otimes \theta),$$

where $u = \sum \gamma(\beta)\alpha$ with $R = \sum \alpha \otimes \beta$.

\Rightarrow invariant for **tangles**.



Notation

$A = (A, \eta, m, \varepsilon, \Delta, \gamma)$: a fin-dim Hopf algebra over a field k ,
with basis $\{e_\alpha\}_\alpha$.

$A^{\text{op}} = (A, \eta, m^{\text{op}}, \varepsilon, \Delta, \gamma^{-1})$: the opposite Hopf algebra of A ,

$(A^{\text{op}})^* = (A^*, \varepsilon^*, \Delta^*, \eta^*, (m^{\text{op}})^*, (\gamma^{-1})^*)$: the dual of A^{op} .

Drinfeld double and Heisenberg double

The Drinfeld double $D(A)$ is a **quasi-triangular Hopf algebra** with the universal R -matrix

$R = \sum_a (1 \otimes e_a) \otimes (e^a \otimes 1) \in D(A)^{\otimes 2}$ satisfying

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12} \in D(A)^{\otimes 3}.$$

The Heisenberg double $H(A)$ is an **algebra with the S-tensor**

$S = \sum_a (1 \otimes e_a) \otimes (e^a \otimes 1) \in H(A)^{\otimes 2}$ satisfying

$$S_{12}S_{13}S_{23} = S_{23}S_{12} \in H(A)^{\otimes 3}.$$

Drinfeld double and Heisenberg double

Set

$$S' = \sum (1 \otimes \tilde{e}_a) \otimes (e^a \otimes 1) \in H(A)^{\text{op}} \otimes H(A),$$

$$S'' = \sum (1 \otimes e_a) \otimes (\tilde{e}^a \otimes 1) \in H(A) \otimes H(A)^{\text{op}},$$

$$\tilde{S} = \sum (1 \otimes \tilde{e}_a) \otimes (\tilde{e}^a \otimes 1) \in H(A)^{\text{op}} \otimes H(A)^{\text{op}},$$

where $\tilde{e}_a = \gamma(e_a)$ and $\tilde{e}^b = (\gamma^*)^{-1}(e^b)$.

Drinfeld double and Heisenberg double

Theorem (Kashaev '97)

We have $\phi: D(A) \rightarrow H(A) \otimes H(A)^{\text{op}}$ such that

$$\phi^{\otimes 2}(R) = S''_{14} S_{13} \tilde{S}_{24} S'_{23}.$$

Drinfeld double and Heisenberg double

$D(A)$: Drinfeld double of A .

We have a ribbon Hopf algebra

$$\mathfrak{R} = D(A)[\theta] / (\theta^2 - u\gamma(u)),$$

where $u = \sum \gamma^*(e^a) \otimes e_a$.

We also consider the algebra

$$\mathcal{H} = (H(A) \otimes H(A)^{\text{op}}) [\bar{\theta}] / (\bar{\theta}^2 - \phi(u\gamma(u))),$$

and extend the embedding $\phi: D(A) \rightarrow H(A) \otimes H(A)^{\text{op}}$ to the map $\bar{\phi}: \mathfrak{R} \rightarrow \mathcal{H}$ by $\bar{\phi}(\theta) = \bar{\theta}$.

Universal quantum invariant and its reconstruction


Universal quantum invariant for tangles in a cube

(1) Choose a diagram

(2) Put labels



(3) Read labels




$$J(C) = \sum \gamma(\alpha)\gamma(\beta')u\theta^{-1} \otimes \alpha'\beta \in \bar{\mathfrak{R}} \otimes \mathfrak{R}.$$

$$(R = \sum \alpha \otimes \beta = \sum \alpha' \otimes \beta')$$

Reconstruction of the universal quantum invariant


(1) Modify diagram

- ▶ Exchange \cup and \cap with \cup and \cap , resp.
- ▶ Duplicate strands
- ▶ Thicken the left strands




Reconstruction of the universal quantum invariant

(2) Put labels



(3) Read the labels




$$J'(C) = (\bar{\theta} \otimes 1)\phi^2(J(C)) \in \bar{\mathcal{H}} \otimes \mathcal{H}.$$


Sketch of proof

$$J: \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \mapsto R$$

$$J': \begin{array}{c} 1 \quad 2 \quad 3 \quad 4 \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \end{array} \mapsto S''_{14} S_{13} \tilde{S}_{24} S'_{23}$$

Sketch of proof

$$J: \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \mapsto R$$



$$J': \begin{array}{c} 1 \quad 2 \quad 3 \quad 4 \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \end{array} \mapsto S''_{14} S_{13} \tilde{S}_{24} S'_{23} = \phi^{\otimes 2}(R)$$


Extension of the universal quantum invariant

- ▶ Colored diagrams
- ▶ Colored moves
- ▶ Invariance of the universal quantum invariant

Colored diagrams


: tangle diagrams obtained from the following parts



We can define the map J' on colored diagrams in a similar way.

Colored moves


- ▶ Colored Pachner (2, 3) moves



Here, the orientation of each strand is arbitrary, and the thickness of each strand with *-mark is arbitrary.

Colored moves


- ▶ Colored $(0, 2)$ moves



Here, the orientation and thickness of each strand are arbitrary.

Colored moves

► Colored symmetry moves



Here, the orientation and thickness of each strand are arbitrary.

Colored moves

► Planer isotopies

$$\text{Wavy line} = | = \text{Bumpy line},$$

$$\text{Cross with yellow square} = \text{Cross with yellow square and strands}, \quad \text{Cross} = \text{Cross with strands}, \quad \text{Cross} = \text{Cross with strands}$$

Here, the orientation and thickness of each strand are arbitrary.

Invariance of the universal quantum invariant

\mathcal{CD} : the set of colored diagrams


\sim_c : the equivalence relation on \mathcal{CD} generated by colored moves.

Theorem (S)

If $\gamma^2 = 1$, then the map J' is an invariant under \sim_c .

Invariance of the universal quantum invariant

\sim'_c : the equivalence relation on \mathcal{CD} generated by colored moves except for



Theorem (S)


The map J' is an invariant under \sim'_c .

3-dimensional descriptions


- ▶ Colored singular triangulations
- ▶ Colored moves
- ▶ v.s. link complements

Colored tetrahedron

: a tetrahedron with an ordering f_1, f_2, f_3, f_4 of its faces




There are eight types of colored tetrahedra:



Colored singular triangulation $\mathcal{C}(Z)$


Define $\mathcal{C}(Z)$ for a colored diagram Z as follows.

(1) Place tetrahedra




Colored singular triangulation $\mathcal{C}(Z)$

(2) Define star-vertices





Colored singular triangulation $\mathcal{C}(Z)$

(2) Attach the tetrahedra




Colored moves

colored Pachner (2,3) move



pentagon relation


J'



v.s. link complements in $S^3 \setminus \{\pm\infty\}$


The octahedral decomposition $\mathcal{O}(D)$:

(1) Place an octahedron at each crossing



v.s. link complements in $S^3 \setminus \{\pm\infty\}$

(2) Attach the octahedra




the boundary of the octahedron

v.s. link complements in $S^3 \setminus \{\pm\infty\}$

Theorem (S)

The octahedral triangulation $\mathcal{O}(D)$ admits a colored ideal triangulation $\mathcal{C}(Z(D))$.

Sketch of the proof



Remarks

- ▶ $\gamma^2 = 1 \Rightarrow J'$ is an inv. of **closed 3-mfd.**
- ▶ (Conj) $\gamma^2 \neq 1 \Rightarrow J'$ is an inv. of **framed 3-mfd.**
- ▶ The colored diagrams form a strict monoidal category and J' is **formulated as a functor.**
- ▶ Hoping to get TQFT if we take $L^2(\mathbb{R})$ as a module of $H(B_q(sl_2))$, **which may give $\text{Vol}(M) + i\text{CS}(M)$.**